

# Synthesis, structure and luminescence of lanthanide containing coordination polymers

*Roel Decadt<sup>1</sup>, Kristof Van Hecke<sup>1</sup> and Rik Van Deun<sup>1</sup>*

*<sup>1</sup> Ghent University, Department of Inorganic and Physical Chemistry, Krijgslaan 281,  
9000 Gent, Belgium*

In this work, a number of lanthanide coordination polymers were synthesized and characterized through X-ray single crystal and powder diffraction and luminescence spectroscopy.

Four new terephthalic acid linked coordination polymers were synthesized, containing praseodymium, neodymium, samarium and europium.[1] The compounds that did not yield crystals suitable for single crystal analysis were Rietveld-refined based on the powder diffraction patterns. In this series of compounds, the lanthanide contraction was perceived through a proportionate distortion of the crystal lattice. The europium-containing coordination polymer exhibited bright luminescence in the visible (red) region of the spectrum.

A neodymium homologue of 2-5-pyridinedicarboxylic acid-based coordination polymers was also synthesized and characterized.[2] The compound showed typical Nd<sup>3+</sup> near infrared luminescence.

[1] X. Guo, G. Zhu, *Inorg. Chem.*, **2006**, 45, 2581-2587.

[2] Huang, *J. Solid State Chem.*, **2008**, 181, 1731-1737.